Administrative Office: 2	211 Park Avenue South • Minneapolis, Minnesota www.entsc.com		ıx (612) 871-2012
We have made arrangements for Dr		to perform your surgery	
at	Hospital on	Your su	rgery is
scheduled for	a.m. / p.m. Please go to the hospital	Admitting Office at	a.m. / p.m.
The surgery being per	rformed is:		
	CHILDREN AND AD	DULTS	
BEFORE SURGER	Y:		
prior to your surgery f	cal is required withindays befor completion of the enclosed examination for ospital on the day of the surgery; do not mail	orm. You must bring the pre-ope	erative physical
Let your primary care	doctor know if you develop a cold or other in	nfection the week of, or before,	your surgery.
Aspirin, or medication	ns containing aspirin (ecotrin, Excedrin), sho	uld be stopped two weeks befor	re surgery.
ncludes ibuprofen, ind Celebrex, Clinoril, Da	Clammatory medications must be stopped on domethacin, naproxen, ketoprofen, ketorolac, ypro, Dolobid, Feldene, Haltran, Indocin, Lo prin, Orudis, Rufen, Ponstel, Tolectin, Torado	, Advil, Aleve, Anaprox, Ansaid dine, Meclomen, Medipren, Mi	d, Cataflam, idol, Motrin,
ovenox, heparin, pers	r another doctor prescribed medication containantine, plavix), please alert our office before the direction of your primary care doctor. As	your surgery. These will need t	to be discontinued

Discontinue black cohash, gingko baloba two weeks before surgery. Discontinue Vitamin E one week before surgery, discontinue supplements containing ephedra or ephedrine 48 hours before surgery.

No alcoholic beverages are permitted within 48 hours before surgery.

Prescribed medications (blood pressure, heart, lung, seizure, etc.) may be taken the morning of surgery with a small sip of water. If you have questions concerning medications, call your primary care doctor. Do not take any other medications without the knowledge of your doctor.

Tylenol, or acetaminophen, does not affect your blood-clotting and may be taken before or after surgery.

Please arrange for someone else to drive you to and from the hospital. Do not plan to drive yourself.

Please contact us immediately if you need to cancel or reschedule.

Upon arriving home after surgery, if the patient has not scheduled post op appointments, call the clinic immediately.

C/3/04 (OVER)

FOOD AND DRINK GUIDELINES BEFORE SURGERY

Ingested Material

Minimum Fasting Period (hours)

Any Solid Food	8
Clear Liquids	3
Breast Milk	4
Infant Formula /Milk/Liquid Feeding Preparations	6

Clear liquids include only water, popsicles, apple juice, soft drinks, Kool-Aid and Jell-O. Clear liquids do not include hard candy, gum or juice with pulp such as orange juice.

If the patient is on chronic medications or if the patient's doctor recommends continuation of specific medication before surgery, you may administer it as scheduled with a sip of clear liquids, if necessary. Do not give medications with pudding, applesauce or other solid foods.

If the patient has any further questions regarding prescribed medications, please call the hospital or surgery center Anesthesia Department so that they can tell you when the last dosage should be given before surgery.

The patient's teeth may be brushed and rinsed with water only.

If the patient has not followed the above guidelines and has had food or water before surgery, please call the Anesthesia Department at the hospital or surgery center where the procedure is being performed.

If these guidelines for adults and children are not followed, the surgery may be cancelled.

Any further medical questions, please feel free to contact the nurse at the office where you were seen.

Upon arriving home after surgery, if the patient has <u>not</u> scheduled post op appointments, call the clinic immediately.

Thank you.

www.entsc.com

(OVER)

2211 Park Avenue South Minneapolis, MN 55404 612-871-1144 Fax 612-871-2012 602 Garden View Med. Bldg. 347 Smith Avenue North St. Paul, MN 55102 651-227-0821 Fax 651-297-6597